Provided for non-commercial research and education use. Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

http://www.elsevier.com/authorsrights

Author's personal copy

Linear Algebra and its Applications 439 (2013) 2309-2329



Contents lists available at SciVerse ScienceDirect

Linear Algebra and its Applications

www.elsevier.com/locate/laa

LINEAR ALGEBRA and Its Applications

An efficient algorithm for maximum entropy extension of block-circulant covariance matrices



F.P. Carli^{a,*}, A. Ferrante^a, M. Pavon^b, G. Picci^a

^a Department of Information Engineering, University of Padova, Italy ^b Department of Mathematics, University of Padova, Italy

ARTICLE INFO

Article history: Received 28 December 2012 Accepted 12 June 2013 Available online 17 July 2013 Submitted by A. Böttcher

MSC: 15A83 15B05 93E12 60G10 49N15 65K10 *Keywords:* Matrix completion Positive definite completion Reciprocal processes Covariance matrices Circulant matrices

ABSTRACT

This paper deals with maximum entropy completion of partially specified block-circulant matrices. Since positive definite symmetric circulants happen to be covariance matrices of stationary periodic processes, in particular of stationary reciprocal processes, this problem has applications in signal processing, in particular to image modeling. In fact it is strictly related to maximum likelihood estimation of bilateral AR-type representations of acausal signals subject to certain conditional independence constraints. The maximum entropy completion problem for block-circulant matrices has recently been solved by the authors, although leaving open the problem of an efficient computation of the solution. In this paper, we provide an efficient algorithm for computing its solution which compares very favorably with existing algorithms designed for positive definite matrix extension problems. The proposed algorithm benefits from the analysis of the relationship between our problem and the band-extension problem for block-Toeplitz matrices also developed in this paper.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Toeplitz matrices

We consider the problem of completing a partially specified block-circulant matrix under the constraint that the completed matrix should be positive definite and block-circulant with an inverse of

0024-3795/\$ – see front matter © 2013 Elsevier Inc. All rights reserved. http://dx.doi.org/10.1016/j.laa.2013.06.014

^{*} Corresponding author. Present address: Department of Electrical and Computer Engineering, University of Minnesota, 200 Union street S.E., Minneapolis, MN 55455, United States. Tel.: +1 612 626 0095.

E-mail addresses: fpcarli@umn.edu (F.P. Carli), augusto@dei.unipd.it (A. Ferrante), pavon@math.unipd.it (M. Pavon), picci@dei.unipd.it (G. Picci).

banded structure. As shown in [1], a block-circulant completion problem of this kind is a crucial tool for the identification of a class of reciprocal processes. These processes [2–4] are a generalization of Markov processes which are particularly useful for modeling random signals which live in a finite region of time or of the space line, for example images. In this paper we consider stationary reciprocal processes for which we refer the reader to [5,6] and references therein. In particular, stationary reciprocal processes of the autoregressive type can be described by linear models involving a banded block-circulant concentration matrix¹ whose non-zero blocks are the (matrix-valued) parameters of the model.

This problem fits in the general framework of covariance extension problems introduced by A.P. Dempster [7] and studied by many authors (see [8–25] and references therein). A key discovery by Dempster is that the inverse of the maximum entropy completion of a partially assigned covariance matrix has zeros exactly in the positions corresponding to the unspecified entries in the given matrix, a property which, from now on, will be referred to as the *Dempster property* (an alternative, concise proof of this statement can for example be found in [26]).

A relevant fact is that, even *when the constraint of a circulant structure is imposed*, the inverse of the maximum entropy completion maintains the Dempster property. This fact has been first noticed in [1] for a banded structure and then proved in complete generality, i.e. for arbitrarily given elements within a block-circulant structure, in [26]. Otherwise stated, this means that the solution of the Maximum Entropy Block-Circulant Extension Problem (CME) and of the Dempster Maximum Entropy Extension Problem (DME) with data consistent with a block-circulant structure, coincide. Note that this property does not hold, for example, for arbitrarily missing elements in a block-Toeplitz structure: if we ask the completion to be Toeplitz, the maximum entropy extension fails to satisfy the Dempster property unless the given data lie on consecutive bands centered along the main diagonal (see [9] and [14] for a general formulation of matrix extension problems in terms of so-called banded-algebra techniques and for a thorough discussion of the so-called band-extension problem for block-Toeplitz matrices). Moreover, the block-Toeplitz band extension problem can be solved by factorization techniques and is essentially a linear problem. This is unfortunately no longer true when a block-circulant structure is imposed to the extension [27].

The main contribution of this paper is to propose a new algorithm for solving the CME problem. A straightforward application of standard optimization algorithms would be too expensive for large sized problems like those we have in mind for, say, applications to image processing. Here we propose a new procedure which rests on duality theory and exploits information on the structure of the problem as well as the circulant structure for computing the solution of the CME.

Since the solutions of the CME and of the DME with circulant-compatible data coincide, methods available in the literature for the DME can, in principle, be employed to compute the solution of CME. In this respect, it has been shown that if the graph associated with the specified entries is chordal [28], the solution of the DME can be expressed in closed form in terms of the principal minors of the covariance matrix, see [16,29,30]. In our problem however the sparsity pattern associated with the given entries is not chordal and the maximum entropy completion has to be computed iteratively. A number of specialized algorithms have been proposed in the graphical models literature; see [7,12, 31,32]. These algorithms deal with the general unstructured setting of Dempster and are not especially tailored to the circulant structure. A detailed comparison of our procedure with the best algorithms available so far is presented in Section 5. We show that the proposed algorithm outperforms the algorithms proposed in the graphical models literature for the solution of the DME, being especially suitable to deal with very large sized instances of the problem.

We shall first relate our work to the solution of the band extension problem for block-Toeplitz matrices and show that the maximum entropy circulant extension approximates arbitrarily closely the block-Toeplitz band extension with the same starting data, when the dimension of the circulant extension becomes large. This result is in the spirit of the relation between stationary Markov and reciprocal processes on an infinite interval established by Levy in [33] and will be useful to provide an efficient initialization for the proposed algorithm. In this context, we shall briefly touch

¹ I.e. the inverse covariance matrix, also known as the precision matrix.

upon feasibility of the CME problem. The feasibility problem for generic blocks size and bandwidth has been addressed in [1] and [26], where a *sufficient* condition on the data for a positive definite *block*-circulant completion to exist has been derived. Here we shall derive a *necessary and sufficient* condition for feasibility of the CME problem valid for the scalar case with unitary bandwidth.

The outline of the paper is as follows. In Section 2 we introduce some notation and state the entropy maximization problem. In Section 3 the relation between the maximum entropy extension for banded Toeplitz and banded circulant matrices is investigated. A necessary and sufficient condition for feasibility is also derived in this section. In Section 4 we describe the proposed procedure for the solution of the CME problem. Section 5 contains a brief review and discussion of some of the most popular methods for the solution of the DME. A comparison of the proposed algorithm and the methods available in the literature is presented in Section 6. Section 7 concludes the paper.

2. Notation and preliminaries

All random variables in this paper, denoted by boldface characters, have zero mean and finite second order moments. It is shown in [1] that a wide-sense stationary \mathbb{R}^m -valued process **y** is stationary on $\{0, 1, ..., N\}$ if and only if its covariance matrix, say Σ_N , has a block-circulant symmetric structure, i.e. Σ_N is of the form

$$\boldsymbol{\Sigma}_{N} = \begin{bmatrix} \Sigma_{0} & \Sigma_{1}^{\top} & \dots & \Sigma_{\tau}^{\top} & \dots & \Sigma_{\tau} & \dots & \Sigma_{1} \\ \Sigma_{1} & \Sigma_{0} & \Sigma_{1}^{\top} & \ddots & \Sigma_{\tau}^{\top} & \dots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & & \ddots & & \Sigma_{\tau} \\ \Sigma_{\tau} & \dots & \Sigma_{1} & \Sigma_{0} & \Sigma_{1}^{\top} & \dots & \ddots & \\ \vdots & \Sigma_{\tau} & \dots & \Sigma_{0} & & \dots & \Sigma_{\tau}^{\top} \\ \vdots & \Sigma_{\tau} & \ddots & & & \vdots \\ \vdots & \ddots & \ddots & & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & & \ddots & \Sigma_{1}^{\top} \\ \Sigma_{1}^{\top} & \dots & \Sigma_{\tau}^{\top} & \dots & \Sigma_{\tau} & & \Sigma_{1} & \Sigma_{0} \end{bmatrix}$$

where the *k*-th block, Σ_k , is given by $\Sigma_k = \mathbb{E}\mathbf{y}(t+k)\mathbf{y}(t)^{\top}$. We refer the reader to [34] for an introduction to circulants; an extension of some relevant results for the *block*-case can be found, for example, in [26]. Here we just recall that the class of block-circulants is closed under sum, product, inverse and transpose. Moreover, all block-circulants are simultaneously diagonalized by the Fourier block-matrix of suitable size (see (9)–(11) below).

The *differential entropy* H(p) of a probability density function p on \mathbb{R}^n is defined by

$$H(p) = -\int_{\mathbb{R}^n} \log(p(x)) p(x) dx.$$
⁽¹⁾

In the case of a zero-mean Gaussian distribution p with covariance matrix Σ_N , it results

$$H(p) = \frac{1}{2} \log(\det \Sigma_N) + \frac{1}{2} n (1 + \log(2\pi)).$$
(2)

Let \mathfrak{S}_N denote the vector space of real symmetric matrices with $N \times N$ square blocks of dimension $m \times m$. Moreover, let \mathbf{U}_N denote the block-circulant shift matrix with $N \times N$ blocks,

$$\mathbf{U}_N = \begin{bmatrix} 0 & I_m & 0 & \dots & 0 \\ 0 & 0 & I_m & \dots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & \dots & I_m \\ I_m & 0 & 0 & \dots & 0 \end{bmatrix},$$

 E_n the $N \times (n+1)$ block matrix

$$E_n = \begin{bmatrix} I_m & 0 & \dots & 0 \\ 0 & I_m & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & I_m \\ 0 & \dots & & 0 \\ \vdots & \ddots & & \vdots \\ 0 & & \dots & 0 \end{bmatrix}$$

and $\mathbf{T}_n \in \mathfrak{S}_{n+1}$ the block-Toeplitz matrix made of the first n + 1, $m \times m$ covariance lags { $\Sigma_0, \Sigma_1, \ldots, \Sigma_n$ },

$$\mathbf{T}_{n} := \begin{bmatrix} \Sigma_{0} & \Sigma_{1}^{\top} & \dots & \Sigma_{n}^{\top} \\ \Sigma_{1} & \Sigma_{0} & \Sigma_{1}^{\top} & & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \Sigma_{1}^{\top} \\ \Sigma_{n} & \dots & \dots & \Sigma_{1} & \Sigma_{0} \end{bmatrix}.$$
(3)

The symmetric block-Toeplitz matrix \mathbf{T}_n is completely specified by its first block-row, so, with obvious notation, it will be also denoted as

$$\mathbf{T}_n = \operatorname{Toepl}(\Sigma_0, \Sigma_1^{\top}, \dots, \Sigma_n^{\top}).$$

Given positive integers N and n such that 2n < N - 1 and a positive definite block-Toeplitz data matrix \mathbf{T}_n , the maximum entropy covariance extension problem for block-circulant matrices (CME) can be stated as follows:

$$\max\{\log \det \boldsymbol{\Sigma}_N \mid \boldsymbol{\Sigma}_N \in \mathfrak{S}_N, \ \boldsymbol{\Sigma}_N > 0\}$$
(4a)

subject to:

$$E_n^{\top} \boldsymbol{\Sigma}_N E_n = \mathbf{T}_n, \tag{4b}$$

$$\mathbf{U}_{N}^{\top}\boldsymbol{\Sigma}_{N}\mathbf{U}_{N}=\boldsymbol{\Sigma}_{N},\tag{4c}$$

where we have exploited the fact that a matrix \mathbf{C}_N with $N \times N$ blocks is block-circulant if and only if it commutes with \mathbf{U}_N , namely if and only if $\mathbf{C}_N = \mathbf{U}_N^{\top} \mathbf{C}_N \mathbf{U}_N$. Problem (4) is a *convex optimization problem* since we are minimizing a strictly convex function on the intersection of a convex cone (minus the zero matrix) with a linear manifold. Existence and uniqueness of the solution of the CME have been proved in [1]. If we do not impose the completion to be block-circulant, we obtain the covariance selection problem studied by A.P. Dempster (DME) in [7].

Notice that, although in problem (4) we are maximizing the entropy functional over zero-mean *Gaussian* densities, we are *not* actually restricting ourselves to the case of Gaussian distributions. Indeed, the Gaussian distribution with (zero mean and) covariance matrix solving (4) maximizes the entropy functional (1) over the larger family of (zero mean) probability densities whose covariance matrix satisfies the boundary conditions (4b), (4c), see [1, Theorem 7.2].

3. Relation with the block-Toeplitz covariance extension problem

In this section, we shall point out a relation between the solutions of the maximum entropy band extension problem for block-circulant and block-Toeplitz matrices.

Let *N* and *n* be positive integers such that n < N, and let \mathbf{T}_n and E_n be the block matrices defined in Section 2. Moreover denote by \mathbf{A}_{N-1} and \mathbf{B}_{N-1} the $(N-1) \times N$ block shift matrices given by

	ΓI_m	0	0		0	0 -			٢0	Im	0	0		ך 0
	0	Im	0		0	0			0	0	Im	0		0
$A_{N-1} =$	0	0	Im		0	0		$B_{N-1} =$	0	0	0	Im	•••	0
	:			•	÷	:	,		:				·	:
	Lo	0	0		Im	0_			Lo	0	0	0	•••	I_m

The maximum entropy band extension problem for block-Toeplitz matrices (TME) can be stated as follows:

$$\max\{\log \det \boldsymbol{\Sigma}_N \mid \boldsymbol{\Sigma}_N \in \mathfrak{S}_N, \ \boldsymbol{\Sigma}_N > 0\}$$
(5a)

subject to:

$$E_n^{\top} \boldsymbol{\Sigma}_N E_n = \mathbf{T}_n, \tag{5b}$$

$$\mathbf{A}_{N-1}\boldsymbol{\Sigma}_{N}\mathbf{A}_{N-1}^{\dagger} = \mathbf{B}_{N-1}\boldsymbol{\Sigma}_{N}\mathbf{B}_{N-1}^{\dagger},\tag{5c}$$

where (5c) constrains Σ_N to be block-Toeplitz by imposing the block-entries on the diagonals to be pairwise equal, and hence constant. This problem has a long history, and was probably the first matrix completion problem studied in the literature [9,14]. The TME problem admits a unique solution. As mentioned in the Introduction, it can be solved by factorization techniques, in fact, by the celebrated Levinson–Whittle algorithm [35] and is essentially a linear problem. Below we shall show that for $N \rightarrow \infty$, the solution of the CME problem can be approximated arbitrarily closely in terms of the solution of the Toeplitz band extension problem. The theorem reads as follows.

Theorem 3.1. Let \mathbf{T}_n be positive definite and let $\{\hat{\Sigma}_k, k = n + 1, n + 2, ...\}$ be the maximum entropy block-Toeplitz extension of $\{\Sigma_0, \Sigma_1, ..., \Sigma_n\}$, that is, the solution of the TME problem (5). Then, for N large enough, the symmetric block-circulant matrix $\boldsymbol{\Sigma}_N^{(c)}$ given by

$$\operatorname{Toepl}(\Sigma_0, \Sigma_1^{\top}, \dots, \Sigma_n^{\top}, \hat{\Sigma}_{n+1}^{\top}, \dots, \hat{\Sigma}_{\frac{N}{2}-1}^{\top}, \hat{\Sigma}_{\frac{N}{2}}^{\top} + \hat{\Sigma}_{\frac{N}{2}}, \hat{\Sigma}_{\frac{N}{2}-1}, \dots, \hat{\Sigma}_{n+1}, \Sigma_n, \dots, \Sigma_1),$$
(6)

for N even, and

$$\operatorname{Toepl}(\Sigma_0, \Sigma_1^{\top}, \dots, \Sigma_n^{\top}, \hat{\Sigma}_{n+1}^{\top}, \dots, \hat{\Sigma}_{\frac{N-1}{2}}^{\top}, \hat{\Sigma}_{\frac{N-1}{2}}, \dots, \hat{\Sigma}_{n+1}, \Sigma_n, \dots, \Sigma_1),$$
(7)

for N odd, is a covariance matrix which for $N \to \infty$ is arbitrarily close to the $mN \times mN$ maximum entropy block-circulant extension of \mathbf{T}_n , i.e. to the solution of the CME problem (4).

Proof. That $\Sigma_N^{(c)}$ is a valid covariance matrix for *N* large enough follows from [1, Theorem 5.1]. It remains to show that $\Sigma_N^{(c)}$ given by (6), (7) tends to the maximum entropy block-circulant extension of \mathbf{T}_n , say Σ_N^o , i.e. that

$$\lim_{N \to \infty} \left\| \boldsymbol{\Sigma}_N^{(c)} - \boldsymbol{\Sigma}_N^o \right\| = 0.$$
(8)

To this aim, we recall that the maximum entropy completion Σ_N^o is the unique completion of the given data whose inverse has the property to be zero in the complementary positions of those assigned [1,26,7]. Thus, (8) holds if and only if for $N \to \infty$ the inverse of $\Sigma_N^{(c)}$ tends to be banded block-circulant

$$\lim_{N\to\infty} \left\| \left(\boldsymbol{\Sigma}_N^{(c)} \right)^{-1} - \left(\boldsymbol{\Sigma}_N^o \right)^{-1} \right\| = 0,$$

i.e. if the off-band blocks of $(\Sigma_N^{(c)})^{-1}$ tend uniformly to zero faster than 1/N.

To show this, recall that $\boldsymbol{\Sigma}_N^{(c)}$ can be block-diagonalized as

$$\boldsymbol{\Sigma}_{N}^{(c)} = \mathbf{V}\boldsymbol{\Psi}_{N}\mathbf{V}^{*} \tag{9}$$

where **V** is the Fourier block-matrix whose (k, l)-th block is

$$V_{kl} := 1/\sqrt{N} \exp[-j2\pi (k-1)(l-1)/N] I_m$$
(10)

and Ψ_N is the block-diagonal matrix

$$\Psi_N := \operatorname{diag}(\Psi_0, \Psi_1, \dots, \Psi_{N-1}), \tag{11}$$

whose diagonal blocks Ψ_{ℓ} are the coefficients of the finite Fourier transform of the first block row of $\Sigma_N^{(c)}$

$$\Psi_{\ell} = \hat{\Sigma}_{0} + e^{j\vartheta_{\ell}}\hat{\Sigma}_{1}^{\top} + (e^{j\vartheta_{\ell}})^{2}\hat{\Sigma}_{2}^{\top} + \dots + (e^{j\vartheta_{\ell}})^{N-2}\hat{\Sigma}_{2} + (e^{j\vartheta_{\ell}})^{N-1}\hat{\Sigma}_{1},$$
(12)

with $\vartheta_{\ell} := -2\pi \ell/N$. Thus in particular

$$\left(\boldsymbol{\Sigma}_{N}^{(c)}\right)^{-1} = \mathbf{V}\boldsymbol{\Psi}_{N}^{-1}\mathbf{V}^{*}$$

where

$$\Psi_N^{-1} := \operatorname{diag}(\Psi_0^{-1}, \Psi_1^{-1}, \dots, \Psi_{N-1}^{-1}).$$

Now, let us consider the block-Toeplitz band extension of the given data \mathbf{T}_n , { $\hat{\Sigma}_k$, k = 0, 1, 2, ...}, and the associated spectral density matrix

$$\Phi(z) := \hat{\Sigma}_0 + \sum_{i=1}^{\infty} \hat{\Sigma}_i z^{-i} + \left(\sum_{i=1}^{\infty} \hat{\Sigma}_i z^{-i}\right)^*.$$
(13)

It is well-known [35] that $\Phi(z)$ can be expressed in factored form as

$$\Phi(z) = \left[L_n(z^{-1})\right]^{-1} \Lambda_n \left[L_n(z^{-1})\right]^{-*}$$
(14)

where $L_n(z^{-1})$ is the *n*-th Levinson–Whittle matrix polynomial associated with the block-Toeplitz matrix \mathbf{T}_n

$$L_n(z^{-1}) = \sum_{k=0}^n A_n(k) z^{-k}$$
(15)

with the $A_n(k)$'s and $\Lambda_n = \Lambda_n^\top > 0$ being the solutions of the Yule–Walker type equation

$$[A_n(0) \quad A_n(1) \quad \dots \quad A_n(n)]\mathbf{T}_n^{\top} = [A_n \quad 0 \quad \dots \quad 0].$$
(16)

Note that $\Phi(z)^{-1} = L_n(z^{-1})^* \Lambda_n^{-1} L_n(z)$ is a Laurent polynomial, that can be written as

$$\Phi(z)^{-1} = M_0 + (M_1 z + M_2 z^2 + \dots + M_n z^n) + (M_1 z + M_2 z^2 + \dots + M_n z^n)^*.$$

Moreover, the Ψ_{ℓ} 's in (12) can be written as²

$$\Psi_{\ell} = \hat{\Sigma}_{0} + e^{j\vartheta_{\ell}}\hat{\Sigma}_{1}^{\top} + \dots + \left(e^{j\vartheta_{\ell}}\right)^{h}\hat{\Sigma}_{h}^{\top} + e^{-j\vartheta_{\ell}}\hat{\Sigma}_{1} + \dots + \left(e^{-j\vartheta_{\ell}}\right)^{h}\hat{\Sigma}_{h}$$
(17)

where

$$h := \begin{cases} \frac{N-1}{2}, & \text{for } N \text{ odd,} \\ N/2, & \text{for } N \text{ even.} \end{cases}$$

² For *N* even $e^{j\vartheta_{\ell}h} = e^{-j\vartheta_{\ell}h} = -1$, so that $(e^{j\vartheta_{\ell}})^h \hat{\Sigma}_h^\top + (e^{-j\vartheta_{\ell}})^h \hat{\Sigma}_h = -(\hat{\Sigma}_h + \hat{\Sigma}_h^\top)$.

2314

Now, comparing expression (17) with (13), we can write

$$\Psi_{\ell} = \Phi\left(e^{j\vartheta_{\ell}}\right) - \left[\Delta\Phi_{N}\left(e^{j\vartheta_{\ell}}\right) + \Delta\Phi_{N}^{*}\left(e^{j\vartheta_{\ell}}\right)\right]$$
(18)

where

$$\Delta \Phi_N(z) := \sum_{i=h+1}^{\infty} \hat{\Sigma}_i z^{-i}.$$

Since the causal part of $\Phi(z)$ is a rational function with poles inside the unit circle,

$$\sup_{l=0,...,N-1} \left\| \Delta \Phi_N \left(e^{j\vartheta_\ell} \right) + \Delta \Phi_N^* \left(e^{j\vartheta_\ell} \right) \right\| \to 0$$

exponentially fast for $N \to \infty$. It follows that, for $N \to \infty$, Ψ_{ℓ}^{-1} tends to $(\Phi(e^{j\vartheta_{\ell}}))^{-1}$, which is given by

$$\left(\Phi\left(e^{j\vartheta_{\ell}}\right)\right)^{-1} = M_0 + M_1 e^{j\vartheta_{\ell}} + \dots + M_n \left(e^{j\vartheta_{\ell}}\right)^n + M_1^{\top} \left(e^{j\vartheta_{\ell}}\right)^{N-1} + \dots + M_n^{\top} \left(e^{j\vartheta_{\ell}}\right)^{N-n}, \quad (19)$$

for every $\ell = 0, 1, ..., N - 1$. In other words, Ψ_{ℓ}^{-1} tends to the finite Fourier transform of a sequence of the form

 $M_0, M_1^{\top}, M_2^{\top}, \dots, M_n^{\top}, 0, \dots, 0, M_n, \dots, M_1,$

i.e. $(\boldsymbol{\Sigma}_N^{(c)})^{-1}$ tends to be banded block-circulant, as claimed. \Box

This result is very much in the spirit of the findings by Levy [33], which establish a relation between stationary Markov and reciprocal processes on an infinite interval and will be used in Section 4 to provide an efficient initialization for the proposed algorithm.

Feasibility of the CME problem has been addressed in [1], where a sufficient condition on the data for a positive definite block-circulant completion to exist has been derived. There is a simple, yet, to the best of our knowledge, still unnoticed, necessary and sufficient condition for the existence of a positive definite circulant completion for scalar (blocks of size 1×1) entries and bandwidth n = 1 which can be derived by combining the results in [17,7,26]. It reads as follows.

Proposition 3.1. Let $N \ge 4$. The partially specified circulant matrix

Г	σ_0	σ_1	?			?	σ_1
	σ_1	σ_0	σ_1	?			?
	?	σ_1	σ_0	σ_1	?		?
	:			•••			
	•			•			
	:				•••		?
	?	•••		?	σ_1	σ_0	σ_1
	σ_1	?			?	σ_1	σ_0

(20)

admits a positive definite circulant completion if and only if $|\sigma_1| < \sigma_0$, for N even, and $\cos(\frac{N-1}{N}\pi)\sigma_0 < \sigma_1 < \sigma_0$, for N odd.

Proof. In [17, Corollary 5] it is shown that the partially specified circulant matrix (20) admits a positive definite (but not necessarily circulant) completion if and only if $|\sigma_1| < \sigma_0$, for *N* even, and $\cos(\frac{N-1}{N}\pi)\sigma_0 < \sigma_1 < \sigma_0$, for *N* odd. On the other hand, Dempster [7] shows that if there is any positive definite symmetric matrix which agrees with the partially specified one in the given positions, then there exists exactly one such a matrix with the additional property that its inverse has zeros in the complementary positions of those specified and this same matrix is the one which maximizes the entropy functional among all the normal models whose covariance matrix agrees with the given data.

2315

But, accordingly to the findings in [26], if the given data are consistent with a circulant structure, the maximum entropy completion is necessarily circulant, which concludes the proof. \Box

Proposition 3.1 provides an explicit condition on the off-diagonal entries for the CME to be feasible. Moreover, for given σ_0 and σ_1 , it states that feasibility depends on the size *N* of the asked completion. This confirms, by a completely independent argument, the findings in [1, Theorem 5.1], where the dependency of feasibility on the completion size *N* has been first noticed (and proved for a generic block-size and bandwidth). A clarifying example, which also makes use of the characterization of the set of the positive definite completions in [26], is presented in Appendix A.

4. A new algorithm for the solution of the CME problem

In this section we shall derive our new algorithm to solve the CME problem. The derivation rests upon duality theory for the CME problem developed in [1, Section VI] and profits by the structure of our CME along with the properties of block-circulant matrices to devise a computationally advantageous procedure for the computation of its solution.

Consider the CME as defined in (4) and define the linear map

$$A: \mathfrak{S}_{n+1} \times \mathfrak{S}_N \to \mathfrak{S}_N$$
$$(\Lambda, \Theta) \mapsto E_n \Lambda E_n^\top + \mathbf{U}_N \Theta \mathbf{U}_N^\top - \Theta$$
(21)

and the set

$$\mathcal{L}_{+} := \left\{ (\Lambda, \Theta) \in (\mathfrak{S}_{n+1} \times \mathfrak{S}_{N}) \mid (\Lambda, \Theta) \in \left(\ker(\Lambda) \right)^{\perp}, \left(E_{n} \Lambda E_{n}^{\top} + \mathbf{U}_{N} \Theta \mathbf{U}_{N}^{\top} - \Theta \right) > 0 \right\}.$$
(22)

 \mathcal{L}_+ is an open, convex subset of $(\ker(A))^{\perp}$. Letting $\langle A, B \rangle := \operatorname{tr} AB^{\top}$, the Lagrangian function is given by

$$L(\boldsymbol{\Sigma}_{N}, \boldsymbol{\Lambda}, \boldsymbol{\Theta}) := -\operatorname{tr} \log \boldsymbol{\Sigma}_{N} + \langle \boldsymbol{\Lambda}, (\boldsymbol{E}_{n}^{\top} \boldsymbol{\Sigma}_{N} \boldsymbol{E}_{n} - \mathbf{T}_{n}) \rangle, + \langle \boldsymbol{\Theta}, (\mathbf{U}_{N}^{\top} \boldsymbol{\Sigma}_{N} \mathbf{U}_{N} - \boldsymbol{\Sigma}_{N}) \rangle$$

$$= -\operatorname{tr} \log \boldsymbol{\Sigma}_{N} + \operatorname{tr} (\boldsymbol{E}_{n} \boldsymbol{\Lambda} \boldsymbol{E}_{n}^{\top} \boldsymbol{\Sigma}_{N}) - \operatorname{tr} (\boldsymbol{\Lambda} \mathbf{T}_{n}) + \operatorname{tr} (\mathbf{U}_{N} \boldsymbol{\Theta} \mathbf{U}_{N}^{\top} \boldsymbol{\Sigma}_{N}) - \operatorname{tr} (\boldsymbol{\Theta} \boldsymbol{\Sigma}_{N})$$

and its first variation (at Σ_N in direction $\delta \Sigma_N \in \mathfrak{S}_N$) is

$$\delta L(\boldsymbol{\Sigma}_N, \Lambda, \Theta; \delta \boldsymbol{\Sigma}_N) = -\operatorname{tr} (\boldsymbol{\Sigma}_N^{-1} \delta \boldsymbol{\Sigma}_N) + \operatorname{tr} (E_n \Lambda E_n^{\top} \delta \boldsymbol{\Sigma}_N) + \operatorname{tr} ((\mathbf{U}_N \Theta \mathbf{U}_N^{\top} - \Theta) \delta \boldsymbol{\Sigma}_N).$$

Thus $\delta L(\boldsymbol{\Sigma}_N, \Lambda, \Theta; \delta \boldsymbol{\Sigma}_N) = 0$, $\forall \delta \boldsymbol{\Sigma}_N \in \mathfrak{S}_N$ if and only if

$$\boldsymbol{\Sigma}_N^{-1} = \boldsymbol{E}_n \boldsymbol{\Lambda} \boldsymbol{E}_n^\top + \boldsymbol{\mathsf{U}}_N \boldsymbol{\varTheta} \boldsymbol{\mathsf{U}}_N^\top - \boldsymbol{\varTheta}.$$

It follows that, for each fixed pair $(\Lambda, \Theta) \in \mathcal{L}_+$, the unique Σ_N^o minimizing the Lagrangian over $\mathfrak{S}_{N,+} := \{\Sigma_N \in \mathfrak{S}_N, \ \Sigma_N > 0\}$ is

$$\boldsymbol{\Sigma}_{N}^{o} = \left(E_{n}\Lambda E_{n}^{\top} + \mathbf{U}_{N}\boldsymbol{\Theta}\mathbf{U}_{N}^{\top} - \boldsymbol{\Theta}\right)^{-1}.$$
(23)

Moreover, computing the Lagrangian at $\Sigma_N = \Sigma_N^o$ results in

$$L(\boldsymbol{\Sigma}_{N}^{o}, \Lambda, \boldsymbol{\Theta}) = -\operatorname{tr}\log((E_{n}\Lambda E_{n}^{\top} + \mathbf{U}_{N}\boldsymbol{\Theta}\mathbf{U}_{N}^{\top} - \boldsymbol{\Theta})^{-1}) + \operatorname{tr}[(E_{n}\Lambda E_{n}^{\top} + \mathbf{U}_{N}\boldsymbol{\Theta}\mathbf{U}_{N}^{\top} - \boldsymbol{\Theta})(E_{n}\Lambda E_{n}^{\top} + \mathbf{U}_{N}\boldsymbol{\Theta}\mathbf{U}_{N}^{\top} - \boldsymbol{\Theta})^{-1}] - \operatorname{tr}(\Lambda \mathbf{T}_{n}) = \operatorname{tr}\log(E_{n}\Lambda E_{n}^{\top} + \mathbf{U}_{N}\boldsymbol{\Theta}\mathbf{U}_{N}^{\top} - \boldsymbol{\Theta}) + \operatorname{tr}I_{mN} - \operatorname{tr}(\Lambda \mathbf{T}_{n}).$$

This is a strictly concave function on \mathcal{L}_+ whose maximization is the *dual problem* of (CME). We can equivalently consider the convex problem

$$\min\{J(\Lambda,\Theta), (\Lambda,\Theta) \in \mathcal{L}_+\},\tag{24}$$

where J is given by

$$J(\Lambda, \Theta) = \operatorname{tr}(\Lambda \mathbf{T}_n) - \operatorname{tr}\log(E_n \Lambda E_n^\top + \mathbf{U}_N \Theta \mathbf{U}_N^\top - \Theta).$$
⁽²⁵⁾

It can be shown [1, Theorem 6.1] that the function J admits a unique minimum point $(\overline{\Lambda}, \overline{\Theta})$ in \mathcal{L}_+ . The gradient of J with respect to Λ is

$$\nabla_{\Lambda} J(\Lambda, \Theta) = -E_n^{\top} \left(E_n \Lambda E_n^{\top} + \mathbf{U}_N \Theta \mathbf{U}_N^{\top} - \Theta \right)^{-1} E_n + \mathbf{T}_n,$$
(26)

Thus the application of whatever first-order iterative method for the minimization of J would involve repeated inversions of the $mN \times mN$ block matrix $(E_n \Lambda E_n^\top + \mathbf{U}_N \Theta \mathbf{U}_N^\top - \Theta)$, which could be a prohibitive task for N large. Nevertheless, by exploiting our knowledge of the problem, we can devise the following alternative. Let $(\bar{\Lambda}, \bar{\Theta})$ be the unique minimum point of the functional J on \mathcal{L}_+ . We know that $(\bar{\Lambda}, \bar{\Theta})$ are such that $\Sigma_N^o = (E_n \bar{\Lambda} E_n^\top + U_N \bar{\Theta} U_N^\top - \bar{\Theta})^{-1}$ is circulant. Thus, one can think of restricting the search for the solution of the optimization problem to the set

$$\left\{ (\Lambda, \Theta) \mid \left(E_n \Lambda E_n^\top + \mathbf{U}_N \Theta \mathbf{U}_N^\top - \Theta \right) \text{ is circulant} \right\}.$$
(27)

If we denote by \mathfrak{C}_N the linear subspace of symmetric, block-circulant matrices, by \mathfrak{C}_N^{\perp} its orthogonal complement, and by $\Pi_{\mathfrak{C}_N}$ and $\Pi_{\mathfrak{C}_N^{\perp}}$ the orthogonal projection on \mathfrak{C}_N and \mathfrak{C}_N^{\perp} , respectively, set (27) can be written as

$$\left\{ (\Lambda, \Theta) \mid \Pi_{\mathfrak{C}_{N}^{\perp}} \left(E_{n} \Lambda E_{n}^{\top} + \mathbf{U}_{N} \Theta \mathbf{U}_{N}^{\top} - \Theta \right) = 0 \right\}.$$

$$(28)$$

We can now exploit the characterization of the matrices belonging to the orthogonal complement of \mathfrak{C}_N in [1, Lemma 6.1], which states that a symmetric matrix M belongs to the orthogonal complement of \mathfrak{C}_N , if and only if, for some $N \in \mathfrak{S}_N$, it can be expressed as $M = \mathbf{U}_N N \mathbf{U}_N^\top - N$. Thus $(\mathbf{U}_N \Theta \mathbf{U}_N^\top - \Theta) \in \mathfrak{C}_N^\perp$ and set (28) can be written as

$$\left\{ (\Lambda, \Theta) \mid \Pi_{\mathfrak{C}_{N}^{\perp}} \left(E_{n} \Lambda E_{n}^{\top} \right) = - \left(\mathbf{U}_{N} \Theta \mathbf{U}_{N}^{\top} - \Theta \right) \right\}.$$
⁽²⁹⁾

If we compute the dual function J on the set (29), we obtain

$$J(\Lambda, \Theta)|_{\{(\Lambda, \Theta)|\Pi_{\mathfrak{C}_{N}^{\perp}}(E_{n}\Lambda E_{n}^{\top})=-(\mathbf{U}_{N}\Theta\mathbf{U}_{N}^{\top}-\Theta)\}}$$

$$= \operatorname{tr}(\Lambda\mathbf{T}_{n}) - \operatorname{tr}\log(E_{n}\Lambda E_{n}^{\top}+\mathbf{U}_{N}\Theta\mathbf{U}_{N}^{\top}-\Theta^{\top})$$

$$= \operatorname{tr}(\Lambda\mathbf{T}_{n}) - \operatorname{tr}\log(E_{n}\Lambda E_{n}^{\top}-\Pi_{\mathfrak{C}_{N}^{\perp}}(E_{n}\Lambda E_{n}^{\top}))$$

$$= \operatorname{tr}(\Lambda\mathbf{T}_{n}) - \operatorname{tr}\log(\Pi_{\mathfrak{C}_{N}}(E_{n}\Lambda E_{n}^{\top}))$$
(30)

where an explicit formula for the orthogonal projection of $E_n \Lambda E_n^{\top}$ on the subspace of symmetric, block-circulant matrices is given by Theorem 7.1 in [1]. In fact, if we denote

$$\Lambda = \begin{bmatrix} \Lambda_{00} & \Lambda_{01} & \dots & \Lambda_{0n} \\ \Lambda_{01}^\top & \Lambda_{11} & \dots & \Lambda_{1n} \\ \vdots & & \ddots & \vdots \\ \Lambda_{0n}^\top & \Lambda_{1n}^\top & \dots & \Lambda_{nn} \end{bmatrix},$$

it can be shown that the orthogonal projection of $E_n \Lambda E_n^{\top}$ onto \mathfrak{C}_N , say Π_{Λ} , is the banded blockcirculant matrix given by

Author's personal copy

F.P. Carli et al. / Linear Algebra and its Applications 439 (2013) 2309-2329

with

$$\Pi_{0} = \frac{1}{N} (\Lambda_{00} + \Lambda_{11} + \dots + \Lambda_{nn}), \tag{31a}$$

$$\Pi_{1} = \frac{1}{N} (\Lambda_{01} + \Lambda_{12} + \dots + \Lambda_{n-1,n})^{\top},$$
(31b)

$$\Pi_n = \frac{1}{N} \Lambda_{0n}^{\top}, \tag{31c}$$

and $\Pi_i = 0$, for all *i* in the interval $n + 1 \le i \le N - n - 1$. Let us denote with \overline{J} the restriction of *J* on (29)

$$\overline{J}(\Lambda) := \operatorname{tr}(\Lambda \mathbf{T}_n) - \operatorname{tr}\log\{\Pi_{\mathfrak{C}_N}(E_n\Lambda E_n^{\top})\}.$$
(32)

We consider the gradient

$$\nabla_{\Lambda} \bar{J}(\Lambda) = -E_n^{\top} \Pi_{\Lambda}^{-1} E_n + \mathbf{T}_n.$$
(33)

Again, the computation of the gradient matrix involves the inversion of an $mN \times mN$ matrix, namely the projection on the subspace of symmetric block-circulant matrices of $E_n \Lambda E_n^{\top}$, Π_{Λ} . Nevertheless, notice that this time the $mN \times mN$ matrix to be inverted is *block-circulant*, which implies that its inverse can be efficiently computed by exploiting the block-diagonalization

$$\Pi_{\Lambda} = \mathbf{V} \boldsymbol{\Omega}_{N} \mathbf{V}^{*}, \tag{34}$$

where **V** is the block-Fourier matrix (10) and Ω_N is the block-diagonal matrix whose diagonal blocks are the coefficients of the finite Fourier transform of the first block row of Π_A . In fact, (34) yields

$$\Pi_{\Lambda}^{-1} = \mathbf{V}\boldsymbol{\Omega}_{N}^{-1}\mathbf{V}^{*},$$

so that the cost of computing Π_{Λ}^{-1} reduces to the cost of singularly inverting the $m \times m$ diagonal blocks of Ω_N and indeed, by exploiting the Hermitian symmetry of the diagonal blocks of Ω_N , to the cost of inverting only the first $\lceil \frac{N+1}{2} \rceil m \times m$ blocks of Ω_N . As a final improvement, notice that due to the final left and right multiplication by E_n^{\top} and E_n , only the first n+1 blocks of Π_{Λ}^{-1} are needed to compute the gradient.

To recap, the proposed procedure reduces the computational cost of each iteration of a generic first-order descent method to $O(m^3)$ flops, in place of the $O(N^3)$ operations per iteration which would have been required by a straightforward application of duality theory.

In the following, we apply a gradient descent method to the optimization of the modified functional \overline{J} . The overall proposed procedure is as follows.

Algorithm 1 Matricial gradient descent algorithm

```
Given a starting point \Lambda \in \text{dom } \overline{J}, \alpha \in (0, 0.5), \beta \in (0, 1)

while \|\nabla_{\Lambda} \overline{J}(\Lambda)\|_{2} > \eta do

\Delta \Lambda := -\nabla_{\Lambda} \overline{J}(\Lambda), t := 1

while \overline{J}(\Lambda + t\Delta\Lambda) > \overline{J}(\Lambda) + \alpha t \operatorname{tr}\{\nabla \overline{J}(\Lambda)^{\top} \Delta\Lambda\} do

t := \beta t

end while

\Lambda := \Lambda + t\Delta\Lambda

end while
```

In the next subsection we provide an efficient initialization for Algorithm 1.

A comparison of the proposed procedure with state of the art algorithms for DME from the literature will be presented in Section 6.

4.1. Algorithm initialization

In the following, we exploit the asymptotic result in Theorem 3.1 to provide a good starting point for the iterative procedure of Algorithm 1. To this aim, recall that the maximum entropy completion of a partially specified block-*Toeplitz* matrix can be computed via the formula

$$\boldsymbol{\Phi}(\boldsymbol{z}) = \left(\boldsymbol{G}^{*}(\boldsymbol{z})\mathbf{T}_{n}^{-1}\tilde{\boldsymbol{B}}\left(\tilde{\boldsymbol{B}}^{*}\mathbf{T}_{n}^{-1}\tilde{\boldsymbol{B}}\right)^{-1}\tilde{\boldsymbol{B}}^{*}\mathbf{T}_{n}^{-1}\boldsymbol{G}(\boldsymbol{z})\right)^{-1}$$
(35)

(see [36] for details), where

$$G(z) = (zI - \tilde{A})^{-1}\tilde{B}$$
(36)

with

$$\tilde{B} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ I \end{bmatrix}, \qquad \tilde{A} = \begin{bmatrix} 0 & I & 0 & \dots & 0 \\ 0 & 0 & I & \dots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & & & I \\ 0 & \dots & \dots & 0 \end{bmatrix}.$$
(37)

It follows that the spectral factor $W(z) := [L_n(z^{-1})]^{-1} \Lambda_n^{\frac{1}{2}}$ has a realization

$$W(z) = C(zI - A)^{-1}B + D$$

with $D = A_n^{\frac{1}{2}}$, $C = -[A_n(n) A_n(n-1) \dots A_n(1)]$ and

$$A = \begin{bmatrix} 0 & I_m & 0 & 0 & \dots & 0 \\ 0 & 0 & I_m & 0 & \dots & 0 \\ \vdots & & \ddots & & \vdots \\ \vdots & & & \ddots & 0 \\ 0 & \dots & \dots & \dots & 0 & I_m \\ -A_n(n) & -A_n(n-1) & \dots & \dots & \dots & -A_n(1) \end{bmatrix}, \qquad B = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ A_n^{\frac{1}{2}} \end{bmatrix}.$$

The positive real part of the maximum entropy spectrum is given by

$$\Phi_{+}(z) = C(zI - A)^{-1}\bar{C}^{\top} + \frac{1}{2}\Sigma_{0}$$
(38)



Fig. 1. CPU time (in seconds) for the matricial gradient descent algorithm with different initializations: identity (dashed line) and via the solution of the associated matrix extension problem for Toeplitz matrices (solid line). The reported times have been computed for N = [10, 20, 30, 40, 50], m = 5 and bandwidth n = 3.

Table 1

CPU time (in seconds) for the matricial gradient descent algorithm with different initializations: identity (on the left) and via the solution of the associated matrix extension problem for Toeplitz matrices (on the right). The reported times have been computed for N = [10, 20, 30, 40, 50], m = 5 and bandwidth n = 3.

Ν	т	Identity		Toeplitz			
		# of itz.	CPU time	# of itz.	CPU time		
10	5	99	0.1455	61	0.0767		
20	5	212	0.4143	65	0.1270		
30	5	322	0.8355	97	0.2504		
40	5	432	1.4233	130	0.4285		
50	5	541	2.1937	163	0.6603		

where $\bar{C}^{\top} = APC^{\top} + BD^{\top}$, with $P = APA^{\top} + BB^{\top}$ and the maximum entropy covariance extension results

$$\hat{\Sigma}_k = C A^{k-1} \bar{C}^\top, \quad k > n.$$

With this extension at hand, we can compute an approximation for the maximum entropy blockcirculant extension as suggested by Theorem 3.1. A good starting point for our algorithm can then be obtained from (31) assuming for Λ a Toeplitz structure.

As an example, we have compared the execution time of the proposed algorithm initialized with the identity matrix and initialized with the solution of the associated matrix extension problem for Toeplitz matrices as described above for blocks of size m = 5, bandwidth n = 3 and N varying between 10 and 50. The computational times are reported in Fig. 1 along with Table 1. The simulation results confirm that the proposed initialization acts effectively to reduce the number of iterations (and thus the computational time) required to reach the minimum.

5. Algorithms for the unstructured covariance selection problem

In this section we introduce and discuss some of the main algorithms in the literature for the positive definite matrix completion problem with the aim of comparison with our newly proposed algorithm.



Fig. 2. Banded pattern of the given entries for the TME problem with N = 8, n = 2, m = 1 (a) and associated graph (b).



Fig. 3. Banded pattern of the given entries for the CME with N = 12, n = 3, m = 1 (a) and associated graph (b). The graph is *not* chordal since, for example, the cycle {1, 4, 7, 10} does not have a chord.

In the literature concerning matrix completion problems, it is common practice to describe the pattern of the specified entries of an $mN \times mN$ partial symmetric matrix $M = (m_{ij})$ by an undirected graph of mN vertices which has an edge joining vertex *i* and vertex *j* if and only if the entry m_{ij} is specified. Since the diagonal entries are all assumed to be specified, we ignore loops at the vertices.

As anticipated in the Introduction, if the graph of the specified entries is *chordal* (i.e., a graph in which every cycle of length greater than three has an edge connecting nonconsecutive nodes, see e.g. [28]), the maximum determinant matrix completion problem admits a closed form solution in terms of the principal minors of the sample covariance matrix (see [16,29,30]). An example of chordal sparsity pattern along with the associated graph is shown in Fig. 2. However, the graph associated with a banded circulant sparsity patterns is not chordal, as it is apparent from the example of Fig. 3(b). Therefore we have to resort to *iterative algorithms*. For the applications we have in mind, we are dealing with vector-valued processes possibly defined on a quite large interval. A straightforward application of standard optimization algorithms is too expensive for problems of such a size, and a number of specialized algorithms have been proposed in the graphical models literature

[7,12,31,32]. In his early work [7], Dempster himself proposed two iterative algorithms which however are very demanding from a computational point of view. Two popular methods are those proposed by T.P. Speed and H.T. Kiiveri in [12], that we now briefly discuss.

Speed and Kiiveri's algorithms. We will denote an undirected graph by $\mathcal{G} = (V, E)$, where V is the vertex set and E the edge set which consists of unordered pairs of distinct vertices. In any undirected graph we say that 2 vertices $u, v \in V$ are *adjacent* if $(u, v) \in E$. For any vertex set $S \subseteq V$, consider the edge set $E(S) \subseteq E$ given by

$$E(S) := \{ (u, v) \in E \mid u, v \in S \}.$$

The graph $\mathcal{G}(S) = (S, E(S))$ is called subgraph of \mathcal{G} induced by S. An induced subgraph $\mathcal{G}(S)$ is *complete* if the vertices in S are pairwise adjacent in \mathcal{G} . A *clique* is a complete subgraph that is not contained within another complete subgraph. Finally, we define the complementary graph of $\mathcal{G} = (V, E)$ as the graph $\tilde{\mathcal{G}}$ with vertex set V and edge set \tilde{E} with the property that $(u, v) \in \tilde{E}$ if and only if $u \neq v$ and $(u, v) \notin E$.

Let \mathcal{I}_b be the set of pairs of indices consistent with a banded, symmetric block-circulant structure of bandwidth *n*, i.e. the set of the (i, j)'s which satisfies the following rules set

for
$$i \in \{1, ..., m\}$$
, $j \in \{i, ..., mN\}$, if $|i - j| \leq m(n + 1) - i \Rightarrow (i, j) \in \mathcal{I}_b$,
if $(i, j) \in \mathcal{I}_b \Rightarrow ((i + m)_{\text{mod } mN}, (j + m)_{\text{mod } mN}) \in \mathcal{I}_b$,
if $(i, j) \in \mathcal{I}_b \Rightarrow (j, i) \in \mathcal{I}_b$

(an example of this structure is shown in Fig. 3(a)). Moreover, we will denote by \mathcal{I}_b^C the complement of \mathcal{I}_b with respect to $\{1, \ldots, mN\} \times \{1, \ldots, mN\}$ and by $\mathcal{G} = (\{1, \ldots, mN\}, \mathcal{I}_b)$ the graph associated with the given entries.

As mentioned in the Introduction, for the class of problems studied by Dempster, the inverse of the unique completion which maximizes the entropy functional has the property to be zero in the complementary positions of those fixed in Σ_N . Thus, a rather natural procedure to compute the solution of the covariance selection problem for block-circulant matrices seems to be the following: iterate maintaining the elements of Σ_N indexed by \mathcal{I}_b at the desired value (i.e. equal to the corresponding elements in the sample covariance matrix) while forcing the elements of Σ_N^{-1} in \mathcal{I}_b^C to zero. To this aim, the following procedure can be devised.

Algorithm 2 First algorithm (Speed and Kiiveri [12])

Compute all the cliques \tilde{c}_t in the complementary graph $\tilde{\mathcal{G}}$, say { \tilde{c}_t , $t = 1,, n_{\tilde{c}_t}$ };
Initialize $\boldsymbol{\Sigma}_{N}^{(0)} = \mathbf{R}_{N};$
while some stopping criterion is satisfied do
for all the cliques \tilde{c}_t in $\tilde{\mathcal{G}}$ do
$\boldsymbol{\Sigma}_{N}^{(t)} = \boldsymbol{\Sigma}_{N}^{(t-1)} + \phi \left(\boldsymbol{\Sigma}_{N}^{(t-1)} ight)$
end for

end for end while

Here $\phi(\boldsymbol{\Sigma}_N^{(t-1)})$ is the $mN \times mN$ zero matrix which equals

$$\{\operatorname{diag}[((\boldsymbol{\Sigma}_{N}^{(t-1)})^{-1})_{\tilde{c}_{t}}]^{-1}\}^{-1} - [((\boldsymbol{\Sigma}_{N}^{(t-1)})^{-1})_{\tilde{c}_{t}}]^{-1}$$

in the positions corresponding to the current clique \tilde{c}_t (given an $mN \times mN$ matrix M and a set $a \subset \{1, \ldots, Nm\}$, M_a denotes the submatrix with entries $\{m_{ij}: i, j \in a\}$). Every cycle consists of as many steps as the cliques in the complementary graph $\tilde{\mathcal{G}}$ (the graph associated to the elements indexed by \mathcal{I}_b^C). At each step, only the elements in $\boldsymbol{\Sigma}_N$ corresponding to the current clique \tilde{c}_t (i.e. only a

subset of the entries indexed by \mathcal{I}_b^C) are modified in such a way to set the elements of $\boldsymbol{\Sigma}_N^{-1}$ in the corresponding positions to the desired zero-value. Throughout the iterations, the elements in $\boldsymbol{\Sigma}_N$ are fixed over \mathcal{I}_b , while the elements of $(\boldsymbol{\Sigma}_N)^{-1}$ vary over \mathcal{I}_b^C .

The role of Σ_N and Σ_N^{-1} can also be swapped, yielding an alternative procedure, which is the analog of iterative proportional scaling (IPS) for contingency tables [37]. Let $\varphi(\Sigma_N^{(t-1)})$ be the $mN \times mN$ zero matrix which equals

$$\left((\mathbf{R}_N)_{c_t}\right)^{-1} - \left(\left(\boldsymbol{\Sigma}_N^{(t-1)}\right)_{c_t}\right)^{-1} \tag{39}$$

in the positions corresponding to the current clique c_t in G (the graph associated with the given entries). The second algorithm reads as follows.

Algorithm 3 Second algorithm (Speed and Kiiveri [12])	
Compute all the cliques c_t in \mathcal{G} , say $\{c_t, t = 1,, n_{c_t}\}$;	
Initialize $\Sigma_N^{(0)} = I_{mN}$;	
while some stopping criterion is satisfied do	
for all the cliques c_t in \mathcal{G} do	
$\left(\boldsymbol{\Sigma}_{N}^{(t)}\right)^{-1} = \left(\boldsymbol{\Sigma}_{N}^{(t-1)}\right)^{-1} + \varphi\left(\boldsymbol{\Sigma}_{N}^{(t-1)}\right)$	(40)
end for end while	

Every cycle consists of as many steps as the cliques in the graph of the specified entries \mathcal{G} . At each step, only the elements in Σ_N^{-1} corresponding to the current clique c_t (i.e. only a subset of the entries indexed by \mathcal{I}_b) are modified in such a way to set the elements of Σ_N in the corresponding positions to the desired value, namely equal to the sample covariance \mathbf{R}_N . Through the iterations the elements in $(\Sigma_N)^{-1}$ are fixed over \mathcal{I}_b^C while the elements of Σ_N vary over \mathcal{I}_b . The choice of which algorithm is to be preferred depends on the application and is very much

The choice of which algorithm is to be preferred depends on the application and is very much dependent on the number and size of the cliques in \mathcal{G} and $\tilde{\mathcal{G}}$. In our setting, the complexity of the graph associated with the given entries depends on the bandwidth *n*. In particular, for a bandwidth *n* not too large with respect to the completion size (which is the case we are interested in) the complexity of the graph associated with the given data \mathcal{G} is far lower than the complexity of its complementary (which, for small *n*, is almost complete), see Fig. 4. The execution time of the two algorithms has been compared for a completion size N = 30 and a bandwidth *n* varying between 2 and 8. The results are shown in Fig. 5 and Table 2. It turns out that for *n* small the second algorithm (which, from now on, will be referred to as IPS) runs faster than the first, and thus has to be preferred.

Covariance selection via chordal embedding. Recently, Dahl, Vanderberghe and Roychowdhury [24] have proposed a new technique to improve the efficiency of the Newton method for the covariance selection problem based on chordal embedding: the given sparsity pattern is embedded in a chordal one for which they provide efficient techniques for computing the gradient and the Hessian. The complexity of the method is dominated by the cost of forming and solving a system of linear equations in which the number of unknowns depends on the number of nonzero entries added in the chordal embedding. For a circulant sparsity pattern, it is easy to check that the number of nonzero elements added in the chordal embedding is quite large. Hence, their method does not seem to be effective for our problem.

6. Comparison of the proposed algorithm and the IPS algorithm

The proposed gradient descent algorithm (GD) applied to the modified dual functional \overline{J} has been compared to the iterative proportional scaling procedure (IPS) by Speed and Kiiveri. Both algorithms are implemented in Matlab. The Bron–Kerbosch algorithm [38] has been employed for finding the

Author's personal copy



Fig. 4. Graph \mathcal{G} associated with the given data (on the left) and its complementary $\tilde{\mathcal{G}}$ (on the right) for N = 20 and bandwidth n = 2, 5, 8.

cliques in the graph for IPS. We recall (see Section 4) that the number of operations per iteration required by our modified gradient descent algorithm is cubic in the block-size *m*, as opposed to the $O([m(N - (n + 1))]^3)$ operations per iteration of the IPS algorithm (see Eqs. (39) and (40)). It follows that for large instances of the CME our newly proposed algorithm is expected to run considerably



Fig. 5. Comparison between the execution time (in seconds) of the first and second algorithm for N = 30, m = 1, $n = \{2, ..., 8\}$.

Execution time of the first and second algorithm for N = 30, m = 1, bandwidth $n = \{2, 2\}$

n	First algorithm		Second algorithm			
	# of cl. (max. cl. size)	CPU time [s]	# of cl. (max. cl. size)	CPU time [s]		
2	4608(10)	9.7877	30(3)	0.4109		
3	2406(7)	4.1515	30(4)	0.1783		
4	1241(6)	1.9419	30(5)	0.3153		
5	706(5)	1.0525	30(6)	0.5535		
6	445(4)	0.6258	30(7)	0.9854		
7	295(3)	0.4145	30(8)	1.7477		
8	175(3)	0.2480	30(9)	3.0665		

faster than the IPS algorithm. The execution times for different completion size N and block size m are plotted in Figs. 6 and 7. The simulation study confirms that our gradient descent algorithm applied to the modified dual functional \overline{J} outperforms the iterative proportional scaling and the gap between the two increases as N increases. Moreover, the gap becomes much more evident as m grows, making the gradient descent algorithm more attractive for applications where the process under observation is vector-valued (m > 1).

7. Conclusions

Table 2

The main contribution of the present paper is an efficient algorithm to solve the maximum entropy band extension problem for block-circulant matrices. This problem has many applications in signal processing since it arises in connection with maximum likelihood estimation of periodic, and in particular quasi-Markov (or reciprocal), processes. Even if matrix completion problems have gained considerable attention in the past (think for example to the covariance extension problem for stationary processes on the integer line, i.e. for Toeplitz matrices), the maximum entropy band extension problem for block-circulant matrices has been addressed for the first time in [1]. The proposed algorithm exploits the circulant structure and relies on the variational analysis brought forth in [1]. An efficient initialization for the proposed algorithm is provided thanks to the established relationship between the solutions of the maximum entropy problem for block-circulant and block-Toeplitz matrices. Further light is also shed on the feasibility issue for the CME problem.

81



Fig. 6. Matricial gradient descent algorithm vs. iterative proportional scaling: CPU time (in seconds) for N = [10, 20, 30, 40, 50], m = 5, bandwidth n = 3.



Fig. 7. Matricial gradient descent algorithm vs. iterative proportional scaling: CPU time (in seconds) for N = [10, 20, 30, 40, 50], m = 10, bandwidth n = 3.

Appendix A. Feasibility of the CME: an example

In Section 3 we have shown that, for given σ_0 and σ_1 , feasibility of the CME depends on the completion size *N*. The following example, aims at clarifying the interplay between feasibility and completion size *N* in the simple case of unitary bandwidth and block-size using the characterization of the set of all positive definite completions derived in [26].

Example A.1. Let $\sigma_0 = 1$, $\sigma_1 = -0.91$. We want to investigate the feasibility of problem (4) for N = 7 and N = 9, i.e. we want to determine if, for N = 7 and N = 9, there exist a positive definite circulant



Fig. A.8. Half-plane $\Psi(w^2) > 0$ and intersection of the half-planes $\Psi(w^0) > 0$ and $\Psi(w^1) > 0$. The intersection of the two regions is empty.

completion for the partially specified matrices

$$\Sigma_7 = \operatorname{Circ}(\sigma_0, \sigma_1, x, y, y, x, \sigma_1), \qquad \Sigma_9 = \operatorname{Circ}(\sigma_0, \sigma_1, x, y, z, z, y, x, \sigma_1),$$

where Circ(a) denotes the circulant symmetric matrix specified by its first row a, and x, y and z denote the unspecified entries. Since

$$\cos\left\{\frac{(N-1)}{N}\pi\right\} = \begin{cases} -0.9010 & \text{for } N = 7, \\ -0.9397 & \text{for } N = 9, \end{cases}$$

by Theorem 3.1, we expect that for N = 7 the problem is unfeasible while for $N \ge 9$ it is expected to become feasible. For N = 7 the set of all positive definite completions is delimited by the intersection of the half-planes identified by the eigenvalues $\Psi(w^k)$, k = 0, ..., 6, of Σ_7

$$\Psi(w^{0}) = -0.82 + 2x + 2y,$$

$$\Psi(w^{1}) = \Psi(w^{6}) = -0.134751 - 0.445042x - 1.80194y$$

$$\Psi(w^{2}) = \Psi(w^{5}) = 1.40499 - 1.80194x + 1.24698y,$$

$$\Psi(w^{3}) = \Psi(w^{4}) = 2.63976 + 1.24698x - 0.445042y$$

(see [26] for details). In Fig. A.8 the intersection Γ of the half-planes $\Psi(w^0) > 0$ and $\Psi(w^1) > 0$ is shown, together with the half-plane $\Psi(w^2) > 0$. The intersection of these two regions is empty. It follows that the intersection of the four half-planes $\Psi(w^k) > 0$, k = 0, ..., 3, is also empty, as claimed. On the other hand, if N = 9, the eigenvalues of Σ_9 are

$$\begin{split} \Psi(w^{0}) &= -0.82 + 2x + 2y + 2z, \\ \Psi(w^{1}) &= \Psi(w^{8}) = -0.394201 + 0.347296x - y - 1.87939z, \\ \Psi(w^{2}) &= \Psi(w^{7}) = 0.68396 - 1.87939x - y + 1.53209z, \\ \Psi(w^{3}) &= \Psi(w^{6}) = 1.91 - x + 2y - z, \\ \Psi(w^{4}) &= \Psi(w^{5}) = 2.71024 + 1.53209x - y + 0.347296z \end{split}$$

and the feasible set is the nonempty region shown in Fig. A.9.



Fig. A.9. Feasible region $\{(x, y, z) | \Sigma_N > 0\}$ for $\Sigma_N = \text{Circ}\{1, -0.91, x, y, z, z, y, x, -0.91\}$.

References

- F. Carli, A. Ferrante, M. Pavon, G. Picci, A maximum entropy solution of the covariance extension problem for reciprocal processes, IEEE Trans. Automat. Control 56 (2011) 1999–2012.
- [2] B. Jamison, Reciprocal processes: The stationary Gaussian case, Ann. Math. Stat. 41 (1970) 1624–1630.
- [3] B. Levy, R. Frezza, A. Krener, Modeling and estimation of discrete-time Gaussian reciprocal processes, IEEE Trans. Automat. Control AC-35 (1990) 1013–1023.
- [4] J. Sand, Four papers in stochastic realization theory, Ph.D. thesis, Dept. of Mathematics, Royal Institute of Technology (KTH), Stockholm, Sweden, 1994.
- [5] B. Levy, A. Ferrante, Characterization of stationary discrete-time Gaussian reciprocal processes over a finite interval, SIAM J. Matrix Anal. Appl. 24 (2002) 334–355.
- [6] A. Ferrante, B. Levy, Canonical form of symplectic matrix pencils, Linear Algebra Appl. 274 (1998) 259–300.
- [7] A. Dempster, Covariance selection, Biometrics 28 (1972) 157–175.
- [8] J. Burg, Maximum entropy spectral analysis, Ph.D. thesis, Dept. of Geophysiscs, Stanford University, Stanford, CA, 1967.
- [9] H. Dym, I. Gohberg, Extension of band matrices with band inverses, Linear Algebra Appl. 36 (1981) 1–24.
- [10] R. Grone, C. Johnson, E. Sa, H. Wolkowicz, Positive definite completions of partial Hermitian matrices, Linear Algebra Appl. 58 (1984) 109–124.
- [11] A. Dembo, C. Mallows, L.A. Shepp, Embedding nonnegative definite Toeplitz matrices in nonnegative definite circulant matrices, with application to covariance estimation, IEEE Trans. Inform. Theory 35 (1989) 1206–1212.
- [12] T. Speed, H. Kiiveri, Gaussian Markov distribution over finite graphs, Ann. Statist. 14 (1986) 138-150.
- [13] C. Johnson, Matrix completion problems: A survey, in: Proc. Sympos. Appl. Math., vol. 40, 1990, pp. 171–198.
- [14] I. Gohberg, S. Goldberg, M. Kaashoek, Classes of Linear Operators, vol. II, Birkhäuser, Boston, 1994.
- [15] I. Gohberg, M. Kaashoek, H. Woerdeman, The band method for positive and strictly contractive extension problems: An alternative version and new applications, Integral Equations Operator Theory 12 (1989) 343–382.
- [16] W. Barrett, C. Johnson, M. Lundquist, Determinantal formulation for matrix completions associated with chordal graphs, Linear Algebra Appl. 121 (1989) 265–289.
- [17] W. Barrett, C. Johnson, P. Tarazaga, The real positive definite completion problem for a simple cycle, Linear Algebra Appl. 192 (1993) 3–31.
- [18] W. Barrett, C. Johnson, R. Loewy, The real positive definite completion problem: Cycle completability, Mem. Amer. Math. Soc. 122 (1996).
- [19] C. Johnson, T. McKee, Structural conditions for cycle completable graphs, Discrete Math. 159 (1996) 155–160.
- [20] M. Laurent, The real positive semidefinite completion problem for series-parallel graphs, Linear Algebra Appl. 252 (1997) 347–366.
- [21] C. Johnson, B. Kroschel, H. Wolkowicz, An interior-point method for approximate positive semidefinite completions, Comput. Optim. Appl. 9 (1998) 175–190.

- [22] W. Glunt, T. Hayden, C. Johnson, P. Tarazaga, Positive definite completions and determinant maximization, Linear Algebra Appl. 288 (1999) 1–10.
- [23] M. Laurent, Polynomial instances of the positive semidefinite and Euclidean distance matrix completion problems, SIAM J. Matrix Anal. Appl. 22 (2001) 874–894.
- [24] J. Dahl, L. Vanderberghe, V. Roychowdhury, Covariance selection for non-chordal graphs via chordal embedding, Optim. Methods Softw. 23 (2008) 501–520.
- [25] A. Ferrante, M. Pavon, Matrix completion à la Dempster by the principle of parsimony, IEEE Trans. Inform. Theory 57 (2011) 3925–3931.
- [26] F. Carli, T. Georgiou, On the covariance completion problem under a circulant structure, IEEE Trans. Automat. Control 56 (2011) 918–922.
- [27] F. Carli, G. Picci, On the factorization approach to band extension of block-circulant matrices, in: Proceedings of the 19th Int. Symposium on the Mathematical Theory of Networks and Systems (MTNS), 2010, pp. 907–914.
- [28] M. Golumbic, Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York, 1980.
- [29] M. Fukuda, M. Kojima, K. Murota, K. Nakata, Exploiting sparsity in semidefinite programming via matrix completion i: General framework, SIAM J. Optim. 11 (2000) 647–674.
- [30] K. Nakata, K. Fujitsawa, M. Fukuda, M. Kojima, K. Murota, Exploiting sparsity in semidefinite programming via matrix completion ii: Implementation and numerical details, Math. Program. Ser. B 95 (2003) 303–327.
- [31] N. Wermut, E. Scheidt, Fitting a covariance selection model to a matrix. Algorithm AS 105, J. R. Stat. Soc. Ser. C. Appl. Stat. 26 (1977) 88–92.
- [32] S. Kullback, Probability densities with given marginals, Ann. Math. Stat. 39 (1968) 1236–1243.
- [33] B. Levy, Regular and reciprocal multivariate stationary Gaussian reciprocal processes over Z are necessarily Markov, J. Math. Syst. Estim. Control 2 (1992) 133–154.
- [34] P. Davis, Circulant Matrices, John Wiley & Sons, 1979.
- [35] P. Whittle, On the fitting of multivariate autoregressions and the approximate spectral factorization of a spectral density matrix, Biometrica 50 (1963) 129–134.
- [36] T. Georgiou, Spectral analysis based on the state covariance: The maximum entropy spectrum and linear fractional parametrization, IEEE Trans. Inform. Theory 52 (2006) 1052–1066.
- [37] S. Haberman, The Analysis of Frequency Data, Univ. Chicago Press, 1974.
- [38] C. Bron, J. Kerbosch, Algorithm 475: Finding all cliques of an undirected graph, Commun. ACM 16 (1973) 575–577.